


- HELAC-NLO - Developments and Applications



Malgorzata Worek
Wuppertal University



Outline

- ❑ General motivation for NLO QCD calculations
- ❑ **HELAC-NLO** in a nutshell
- ❑ Applications: 2 \rightarrow 4 processes:
 - ttbb
 - ttjj
 - WWbb
- ❑ Summary & Outlook

HELAC-NLO Group:

*G. Bevilacqua (INP Athens)
M. Czakon (RWTH Aachen)
M.V. Garzelli (Granada Uní.)
A. van Hameren (INP Kraków)
I. Malamos (Nijmegen Uní.)
C. G. Papadopoulos (INP Athens)
R. Pittau (Granada Uní.)
M. Worek (Wuppertal Uní.)*

Contributors:

*A. Kanaki
A. Cafarella
P. Draggiotis (Valencia Uní.)
G. Ossola (New York City College of Technology)*

Introduction


- ❑ 8-10 partons in the final state @ LO, well separated to avoid divergences
- ❑ On the market automatic parton level tools which are completely self contained
- ❑ Provide amplitudes and integrators on their own
- ❑ Standard Model and beyond tools @ tree level (just few examples)
 - ALPGEN, AMEGIC++/SHERPA, COMIX/SHERPA, **HELAC-PHEGAS**, MADGRAPH/MADEVENT, O'MEGA/WHIZARD, ...
- ❑ General purpose Monte Carlo programs (parton shower, hadronization, multiple interactions, hadrons decays, etc.)
 - HERWIG, HERWIG++, PYTHIA 6.4, PYTHIA 8.1, SHERPA, ...
- ❑ High sensitivity to unphysical input scales, to improve accuracy of prediction higher order calculations are needed

Motivation for $N\mathcal{L}O$

- ❑ Stabilizing the scale in the QCD input parameters most notably the strong coupling constant and PDFs
- ❑ Normalization and shape of distributions first known at NLO
- ❑ Many scale processes: $V + \text{jets}$, $VV + \text{jets}$, $t\bar{t}H$, $t\bar{t} + \text{jets}$, $n\text{jets}$...
- ❑ Sometimes dynamical scales seem to work better for some observables
- ❑ How do we know which scale to choose ?
- ❑ Improved description of jets




Jets: $\mathcal{L}O$



$N\mathcal{L}O$



Parton Shower



Hadron Level

Structure of \mathcal{NLO} Calculations

$$\begin{aligned}\sigma^{NLO} &= \int_m d\sigma^B + \int_{m+1} d\sigma^R - \int_{m+1} d\sigma^A + \int_{m+1} d\sigma^A + \int_m d\sigma^V \\ &\hookrightarrow \int d\sigma^B + \int_{m+1} [d\sigma^R - d\sigma^D] + \int_m [d\sigma^V + d\sigma^I + d\sigma^{KP}]\end{aligned}$$

□ Our strategy in a few words

- make it fully numeric
- make it fully automatic
- „montecarlize“ everything for speed

Virtual Corrections

- Decompose the amplitude into a basis of scalar integrals

$$\mathcal{A} = \sum d_{i_1 i_2 i_3 i_4} \text{ (box) } + \sum c_{i_1 i_2 i_3} \text{ (triangle) } + \sum b_{i_1 i_2} \text{ (bubble) } + \sum a_{i_1} \text{ (self-energy) } + R$$

$$\mathcal{A} = \sum_{I \subset \{0,1,\dots,m-1\}} \int \frac{\mu^{(4-d)d^d q}}{(2\pi)^d} \frac{\bar{N}_I(\bar{q})}{\prod_{i \in I} \bar{D}_i(\bar{q})}$$

- Three main building blocks are needed
 - Evaluation of numerator function $N(q)$
 - Determination of coefficients via reduction method
 - Evaluation of scalar functions via **ONELOOP**

van Hameren, '10

Virtual Corrections

- Reduction at integrand level – **OPP** method implemented in **CUTTOOLS**

- Computing numerator functions for specific values of loop momentum that are solutions of equations

$$\mathcal{D}_i(q) = 0 \quad \text{for } i = 0, \dots, M-1$$

- It is customary to refer to these equations as quadruple ($M = 4$), triple ($M = 3$), double ($M = 2$) and single ($M = 1$) cuts

$$\begin{aligned} N(q) = & \sum_{i_0 < i_1 < i_2 < i_3}^{m-1} \left[d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] \prod_{i \neq i_0, i_1, i_2, i_3}^{m-1} D_i \\ & + \sum_{i_0 < i_1 < i_2}^{m-1} [c(i_0 i_1 i_2) + \tilde{c}(q; i_0 i_1 i_2)] \prod_{i \neq i_0, i_1, i_2}^{m-1} D_i \\ & + \sum_{i_0 < i_1}^{m-1} [b(i_0 i_1) + \tilde{b}(q; i_0 i_1)] \prod_{i \neq i_0, i_1}^{m-1} D_i \\ & + \sum_{i_0}^{m-1} [a(i_0) + \tilde{a}(q; i_0)] \prod_{i \neq i_0}^{m-1} D_i \\ & + \tilde{P}(q) \prod_i^{m-1} D_i. \end{aligned}$$

Virtual Corrections

- ❑ Compute the rational terms $R = R_1 + R_2$
- ❑ R_1 originates from ϵ dependence of denominators

$$D_i \rightarrow \bar{D}_i - \tilde{q}^2$$

- Computed within the framework of OPP reduction

Ossola,, Papadopoulos, Pittau '07, '08

- ❑ R_2 originates from ϵ dependence of numerators


$$\bar{q} = q + \tilde{q} \quad \bar{\gamma}_\mu = \gamma_\mu + \tilde{\gamma}_\mu \quad \bar{g}^{\mu\nu} = g^{\mu\nu} + \tilde{g}^{\mu\nu}$$

- Computed with effective tree-level Feynman rules

*Draggiotis, Garzellí, Papadopoulos, Pittau '09
Garzellí, Malamos, Pittau '09*


Virtual Corrections

- In case of ttbb final state the integrand has the form

$$\mathcal{A}(q) = \sum \underbrace{\frac{N_i^{(6)}(q)}{\bar{D}_{i_0} \bar{D}_{i_1} \cdots \bar{D}_{i_5}}}_{\text{6 blobs}} + \underbrace{\frac{N_i^{(5)}(q)}{\bar{D}_{i_0} \bar{D}_{i_1} \cdots \bar{D}_{i_4}}}_{\text{5 blobs}} + \underbrace{\frac{N_i^{(4)}(q)}{\bar{D}_{i_0} \bar{D}_{i_1} \cdots \bar{D}_{i_3}}}_{\text{4 blobs}} + \underbrace{\frac{N_i^{(3)}(q)}{\bar{D}_{i_0} \bar{D}_{i_1} \bar{D}_{i_2}}}_{\text{3 blobs}} + \cdots$$


- **HELAC-1LOOP** evaluates numerically the numerators $N_i^{(6)}(q)$, $N_i^{(5)}(q)$, ...
- with the values of the loop momentum q provided by **CUTTOOLS**
 - Generates all partitions of 6, 5, 4, ... blobs attached to the loop and checks all possible flavors (colors) that can run inside
 - Hard cuts the loop (q is fixed) to get $n+2$ tree-like process

van Hameren, Papadopoulos, Pittau '09
Bevilacqua, Czakon, Papadopoulos, Pittau, Worek '09





Real Emission

- ❑ **HELAC-DIPOLES** - complete, public, automatic Catani-Seymour dipoles
- ❑ Phase space integration of subtracted real radiation and integrated dipoles in both massless and massive cases
- ❑ Extended for arbitrary polarizations
 - Monte Carlo over polarization states of external particles
 - Monte Carlo over color
- ❑ Phase space restriction on the dipoles phase space
 - Less dipole subtraction terms per event
 - Increased numerical stability
 - Reduced missed binning problem
 - Large cancellations between subtracted real radiation and integrated dipoles

Czakon, Papadopoulos, Worek '09

Real Emission

□ ttbb production @ LHC as an example




$\alpha_{\max} \in (0, 1]$

Bevilacqua, Czakon, Papadopoulos, Pittau, Worek '09


Subtracted real emission
K + P operators
J operators
Full result

Cutoff independence !!!



Evaluation of loop
numerators $N(q)$ and R_2

Reduction of tensor integrals,
OPP coefficients and R_1



Catani-Seymour dipole subtraction for
massless and massive cases

Evaluation of scalar integrals

Applications

$pp \rightarrow tt\bar{b}\bar{b} \text{ \& } pp \rightarrow tt\bar{j}\bar{j}$

$pp \rightarrow t\bar{t}b\bar{b}$ @ LHC

- Integrated cross sections and scale dependence *Per mille level agreement !*

Process	$\sigma_{[23, 24]}^{\text{LO}}$ [fb]	σ^{LO} [fb]	$\sigma_{[23, 24]}^{\text{NLO}}$ [fb]	$\sigma_{\alpha_{\text{max}}=1}^{\text{NLO}}$ [fb]	$\sigma_{\alpha_{\text{max}}=0.01}^{\text{NLO}}$ [fb]
$q\bar{q} \rightarrow t\bar{t}b\bar{b}$	85.522(26)	85.489(46)	87.698(56)	87.545(91)	87.581(134)
$pp \rightarrow t\bar{t}b\bar{b}$	1488.8(1.2)	1489.2(0.9)	2638(6)	2642(3)	2636(3)

$\xi \cdot m_t$	$1/8 \cdot m_t$	$1/2 \cdot m_t$	$1 \cdot m_t$	$2 \cdot m_t$	$8 \cdot m_t$
σ^{LO} [fb]	8885(36)	2526(10)	1489.2(0.9)	923.4(3.8)	388.8(1.4)
σ^{NLO} [fb]	4213(65)	3498(11)	2636(3)	1933.0(3.8)	1044.7(1.7)

$$\sigma_{\text{LO}} = (1489.2 \pm 0.9) \text{ fb}$$

$$\sigma_{\text{NLO}} = (2636 \pm 3) \text{ fb}$$

Scale dependence reduced:

70% @ LO down to **33% @ NLO**

K factor of **K = 1.77**


for quarks initial states only **K = 1.03**

With jet veto of 50 GeV **K = 1.20**

*Bevilacqua, Czakon, Papadopoulos, Pittau, Worek '09
Bredenstein, Denner, Dittmaier, Pozzorini '08, '09*

$pp \rightarrow t\bar{t}b\bar{b}$ @ LHC


Scale dependence graphically



Scale dependence at NLO decomposed into contribution of *Virtual Corrections* & *Real Radiation*

Bevilacqua, Czakon, Papadopoulos, Pittau, Worek '09

Varying scale up or down by a factor two changes cross section by
70% @ LO and by **33% @ NLO**




$pp \rightarrow t\bar{t}b\bar{b}$ @ LHC

- Differential cross sections
- b-jet pair kinematics
 - Invariant mass
 - Transverse momentum
 - Rapidity distribution
- single b-jet kinematics
 - Transverse momentum

\mathcal{LO} & \mathcal{NLO}

- Relatively small variation compared to the size but shape change important



Bevilacqua, Czakon, Papadopoulos, Pittau, Worek '09




$pp \rightarrow t\bar{t}b\bar{b} @ \mathcal{LHC}$

□ Broad study

- *Cross section in fb*
- *Dynamic scale*
- *$m_{b\bar{b}}$ distribution*
- *K-factor*

Bredenstein, Denner, Dittmaier, Pozzorini '10

Setup	$m_{b\bar{b},\text{cut}}$	$p_{T,b\bar{b},\text{cut}}$	$p_{\text{jet,veto}}$	$p_{T,b,\text{cut}}$	$y_{b,\text{cut}}$	σ_{LO}	σ_{NLO}	K
I	100	-	-	20	2.5	786.3(2) ^{+78%} _{-41%}	978(3) ^{+13%} _{-21%}	1.24
II	-	200	-	20	2.5	451.8(2) ^{+79%} _{-41%}	592(4) ^{+13%} _{-22%}	1.31
III	100	-	100	20	2.5	786.1(6) ^{+78%} _{-41%}	700(3) ^{+0.4%} _{-19%}	0.89
IV	100	-	-	50	2.5	419.4(1) ^{+77%} _{-40%}	526(2) ^{+13%} _{-21%}	1.25




$pp \rightarrow t\bar{t}jj$ @ LHC

□ Scale dependence & integrated cross sections

Bevilacqua, Czakon, Papadopoulos, Worek '10

Process	σ^{LO} [pb]	Contribution
$pp \rightarrow t\bar{t}jj$	120.17(8)	100%
$qg \rightarrow t\bar{t}qg$	56.59(5)	47.1%
$gg \rightarrow t\bar{t}gg$	52.70(6)	43.8%
$qq' \rightarrow t\bar{t}qq', q\bar{q} \rightarrow t\bar{t}q'\bar{q}'$	7.475(8)	6.2%
$gg \rightarrow t\bar{t}q\bar{q}$	1.981(3)	1.6%
$q\bar{q} \rightarrow t\bar{t}gg$	1.429(1)	1.2%



$$\sigma_{\text{LO}} = (120.17 \pm 0.08) \text{ pb}$$

$$\sigma_{\text{NLO}} = (106.94 \pm 0.17) \text{ pb}$$

$$\sigma_{\text{NLO}}^{\text{veto}} = (76.58 \pm 0.17) \text{ pb}$$

Scale dependence reduced:

72% @ LO down to **13% @ NLO**

54% @ NLO with **jet veto** of 50 GeV

K factor of **K = 0.89** (**K = 0.64**)

Negative shift of **11%** (**36%**)




$pp \rightarrow ttjj$ @ LHC

□ Differential cross section

LO & NLO

➤ m_{jj} size of the corrections transmitted to distributions for low p_T , shapes change for high p_T

➤ p_T of 1st hardest & 2nd hardest jet (ordered in p_T) altered shapes up to 39% & 28% in tails



Applications


$$pp \text{ (} p\bar{p} \text{)} \rightarrow wwbb$$

$pp (p\bar{p}) \rightarrow wwbb$

- ❑ Complete off-shell effects @ NLO
- ❑ Double-, single- and non-resonant top contributions of the order $O(\alpha_s^3 \alpha^4)$
- ❑ Complex-mass scheme for unstable top
- ❑ W gauge bosons are treated off-shell

$$pp(p\bar{p}) \rightarrow e^+ \nu_e \mu^- \nu_\mu b\bar{b} + X$$

- ❑ Sum over helicities and color via MC
- ❑ LO + V obtained by reweighting of tree level unweighted events
- ❑ Dipole channels for subtracted real part
- ❑ Check of Ward identity for virtual part
- ❑ Cancellation of divergences
- ❑ α_{\max} independence test for real part



$$pp \ (p\bar{p}) \rightarrow wwbb$$

□ Integrated cross sections for inclusive cuts

- $p_T(b) > 20$, GeV, $p_T(l) > 20$ GeV, $p_T(\text{miss}) > 30$ GeV
- $|y(b)| < 4.5$, $|y(l)| < 2.5$, $\Delta R(jj) > 0.4$, $\Delta R(jl) > 0.4$

TeVatron	σ_{LO} [fb]	$\sigma_{\text{NLO}}^{\alpha_{\text{max}}=1}$ [fb]	$\sigma_{\text{NLO}}^{\alpha_{\text{max}}=0.01}$ [fb]
<i>anti-k_T</i>	34.922 ± 0.014	35.705 ± 0.047	35.697 ± 0.049
<i>k_T</i>	34.922 ± 0.014	35.727 ± 0.047	35.723 ± 0.049
C/A	34.922 ± 0.014	35.724 ± 0.047	35.746 ± 0.050

LHC	σ_{LO} [fb]	$\sigma_{\text{NLO}}^{\alpha_{\text{max}}=1}$ [fb]	$\sigma_{\text{NLO}}^{\alpha_{\text{max}}=0.01}$ [fb]
<i>anti-k_T</i>	550.54 ± 0.18	808.46 ± 0.98	808.29 ± 1.04
<i>k_T</i>	550.54 ± 0.18	808.67 ± 0.97	808.86 ± 1.03
C/A	550.54 ± 0.18	808.74 ± 0.97	808.28 ± 1.03



α_{max}
independence test



$pp (p\bar{p}) \rightarrow wwbb$



□ LO & NLO scale dependence

TeVatron

$$\sigma_{LO} = 34.922^{+40\%}_{-26\%} \text{ fb}$$

$$\sigma_{NLO} = 35.727^{+4\%}_{-8\%} \text{ fb}$$

$$K = 1.023$$





LHC

$$\sigma_{LO} = 550.538^{+37\%}_{-25\%} \text{ fb}$$






$$\sigma_{NLO} = 808.665^{+4\%}_{-9\%} \text{ fb}$$

$$K = 1.47$$



$p\bar{p} \rightarrow WWb\bar{b}$






TeVatron



- ☐ Differential cross section
- ☐ Fixed scale m_t
- ☐ Corrections to p_T & $m_{t\bar{t}}$ moderate
- ☐ Do not simply rescale LO shapes
- ☐ Distortions **15% - 80%**
- ☐ Corrections to angular distributions positive and moderate **5% - 10%**




$pp \rightarrow WWb\bar{b}$

LHC



- ☐ Differential cross section
- ☐ Fixed scale m_t
- ☐ Corrections positive and large
50% - 60 %
- ☐ Relatively constant
- ☐ Exception H_T distorted up to **80%**

$$p\bar{p} \rightarrow wwbb$$



$$A_{FB}^t = 0.051$$

$$A_{FB}^b = 0.033$$

$$A_{FB}^{\ell^+} = 0.034$$


□ Asymmetry @ TeVatron

$$A_{FB}^t = \frac{\int_{y>0} N_t(y) - \int_{y<0} N_t(y)}{\int_{y>0} N_t(y) + \int_{y<0} N_t(y)}$$


$$A_{FB}^{\bar{t}} = -A_{FB}^t$$

$pp (p\bar{p}) \rightarrow WWbb$

TeVatron



LHC



- ☐ Size of the non-factorizable corrections
- ☐ Full result versus narrow width approximation
- ☐ Rescaling coupling tWb by some large factors
- ☐ **+1.0%** TeVatron and **-1.2%** LHC for inclusive cuts
- ☐ Dependence of NLO cross section and individual contributions on rescaling parameter ζ

$$\Gamma_{rescaled} = \zeta \Gamma_t$$

$$\Delta\sigma_i(\zeta) = (\sigma_i(\zeta) - \sigma_i(\zeta = 1)) / \sigma_T(\zeta = 1)$$

Bevilacqua, Czakon, van Hameren, Papadopoulos, Worek '11

Summary & Outlook

❑ **ttbb, WWbb** completed by two groups, *Permille level agreement in both cases!*

❑ **ttjj**



See Stefan Kallweit talk on Saturday

❑ **HELAC-NLO**

- Complete tool at NLO built around **HELAC-PHEGAS:**
HELAC-1LOOP, CUTTOOLS, ONELOOP & HELAC-DIPOLES
- Much wider study for ttjj: variation of the center of mass energy, jet algorithms, cone size in jet algorithm, transverse momentum cuts, ...
- Other processes from NLO Wishlist under attack
- Constant improvements in speed and functionality

<http://helac-phegas.web.cern.ch/helac-phegas/>